

Phenolipids: Novel Phenolics Enriched Lecithin for Functional and Pharmaceutical Applications

Ramadan, M. F. ^{*1/2}, Rohn S. ², Mörsel J.-T. ³, Kroh L.W. ²

* Dr. M. F. Ramadan, ¹Agricultural Biochemistry Department, Faculty of Agriculture, Zagazig University, Egypt. ²Technische Universität Berlin, Institut für Lebensmitteltechnologie und Lebensmittelchemie, Fachgebiet Lebensmittelanalytik, TIB 4/3-1, Gustav-Meyer-Allee 25, D-13355 Berlin. ³UBF Untersuchungs-Beratungs-und Forschungslaboratorium GmbH, Altlandsberg, Berlin.

Phenolipids results from the reaction of phospholipids with the selected phenolics in a nonpolar solvent. They are lipophilic substances freely soluble in nonpolar solvents (in which the hydrophilic moiety was not), and moderately soluble in fats. Liposomes, unlike Phenolipids, are formed by mixing water-soluble substances with phospholipids without forming chemical bonds. This difference results in Phenolipids being much better absorbed than liposomes or individual phenolic compounds. Preparation of Phenolipids is recently described by complexing quercetin with soy lecithin [1]. Phenolipids exhibited novel antioxidant properties in a triolein model system stronger than individual lecithin or quercetin.

The goal of this work is to optimize preparation of different structured Phenolipids (phenolics bound or enriched polar lipids). Structure elucidation and antioxidant properties of new Phenolipids are under study. Phenolipids are anticipated to play a vital role in efficient herbal drug delivery of a broad spectrum of protective phytochemicals. After selection of potential phytochemicals from medicinal plants, Phenolipids can be developed for various therapeutic uses like cardiovascular, anti-inflammatory and anticancer activities. Moreover, Phenolipids are anticipated to show their potential in cosmetics as anti-skin ageing agents and for the use of other nonpathogenic skin conditions.

[1] Ramadan MF (2008) Food Science and Technology-LWT 41: 581-587.